

2e. Highlights on Beeja Nirmana - Oogenesis, Spermatogenesis

Bīja-Nirmāṇa (Gamete Formation): Oogenesis & Spermatogenesis — Ayurvedic-Modern Synthesis

Learning Objectives

- Define **Bīja-nirmāṇa** from an Āyurvedic lens and correlate it with modern **oogenesis** and **spermatogenesis**.
- Map **doṣa-dhātu-srotas** principles to the endocrine-cellular events that produce **Artava (female gamete/menstrual element)** and **Śukra (male gamete/seminal element)**.
- Recall high-yield timelines, cell stages, regulatory hormones, and applied aspects relevant to infertility work-ups and preconception care.
- Write exam-ready answers with precise terms, clean tables, and one authenticated classical śloka.

1) Bīja-nirmāṇa in Āyurveda: Core Frame

In the classical doctrine of reproduction, **Bīja** is the seed element of each parent (**Śukra** for the male; **Artava/Rajas** for the female). Healthy progeny (**sūrya-saṃhanana, sātmya, sattva**) depends upon **Rtu-Kṣetra-Ambu-Bīja** (proper time, healthy uterine field, nutritive milieu, and sound gametes). The **channels** (srotas) that conduct and mature these substances are central:

“स्रोतांसि खलु परिणाममापद्यमानानां धातूनामभिवाहीनि भवन्ति ।”

Srotāṃsi khalu pariṇāmam āpadyamānānāṃ dhātūnām abhivāhīni bhavanti.

— **Caraka Saṃhitā, Vimāna Sthāna 5/3**

This verse anchors the **process view**: as dhātus transform, srotas **carry and condition** the evolving material—here, up to the emergence of viable **bīja**.

Link to specific srotas (for quick recall in answers):

- Ārtavavaha srotas (female)**: mūla—**Garbhāśaya + Raktavāhīni dhamanīs** (uterus + vascular network).
 - Śukravaha srotas (male)**: mūla—traditionally described around **Vṛṣaṇa (testes) and śukra-vaha pathways** (seminiferous-epididymal axis in modern terms).
- (Use *Caraka Vimāna-sthāna 5* for srotas authority; write mūla succinctly in your answers.)


2) Oogenesis (Artava-nirmāṇa) — the female bīja

1. Definition

Oogenesis is the process of formation and maturation of female gametes (ova) from primordial germ cells (oogonia) in the ovaries.

2. Stages of Oogenesis

Oogenesis - Stepwise Process



1. Multiplication Phase (Fetal life)

- Primordial germ cells → Oogonia.
- Oogonia multiply by **mitosis** until about 5–7 million are formed.
- By the **5th month of intrauterine life**, oogonia enter meiosis I and become **primary oocytes**.
- At birth: only ~1–2 million primary oocytes remain; at puberty: ~300,000–400,000.

2. Growth Phase

- Primary oocytes remain arrested in **prophase I (diplotene stage)** until puberty (meiotic arrest).
- Each is surrounded by follicular cells → forming **primordial follicles**.
- Oocyte enlarges, cytoplasm accumulates, and a **zona pellucida** forms.

3. Maturation Phase (Puberty to Menopause)

- During each menstrual cycle, under influence of **FSH (follicle-stimulating hormone)**, a few follicles resume meiosis.
- Primary oocyte completes meiosis I → forms:
 - **Secondary oocyte** (large cell, haploid).
 - **First polar body** (small, degenerates).
- Secondary oocyte begins meiosis II but **arrests at metaphase II**.
- Ovulation releases the secondary oocyte.
- Meiosis II completes only if fertilization occurs, forming:
 - **Ovum (mature egg)**.
 - **Second polar body**.

3. Final Outcome

- From one primary oocyte: **only one ovum** is formed (plus 2–3 polar bodies).
- Process is **discontinuous, prolonged, and unequal**.

4. Hormonal Regulation

- **FSH** → follicular growth, oocyte maturation.
- **LH surge** → triggers ovulation, completion of meiosis I.
- **Estrogen & Progesterone** → prepare uterus for implantation.

5. Ayurvedic Insight

- Oogenesis relates to **Beeja Nirmana (female seed)** and **Artava Dhatu** formation.
- Classical texts describe **Rtu-kala** (fertile period) when matured artava is ready for conception.

2.1 Big-picture timeline

Life stage	Event	Key numbers / facts
Fetal life (weeks 6-20)	Oogonia proliferate (mitosis) → enter meiosis I → Primary oocytes	Peak ~ 6-7 million ; all oocytes formed before birth
Late fetal → birth	Atresia begins; all oocytes arrest in prophase I (dictyate)	At birth ~ 1-2 million remain
Puberty	Cyclic recruitment per cohort; most undergo atresia	~ 300-500 thousand at menarche
Each cycle	One dominant follicle completes meiosis I just before ovulation → Secondary oocyte + first polar body ; arrests in metaphase II	Ovulation releases secondary oocyte with cumulus
Fertilization	Sperm entry triggers completion of meiosis II → Ovum + second polar body	~ 400-500 ovulations over reproductive life

Mnemonic: “Made once, paused twice” — made **once** (fetal), **paused** at prophase I and **paused** again at metaphase II.

2.2 Follicular architecture & endocrine control

- **Follicle stages:** Primordial → Primary (granulosa) → Secondary (theca forms) → Antral (Graafian).
- **Two-cell, two-gonadotropin model:**
 - **Theca** cells (LH) → androgens → diffuse to **granulosa**.
 - **Granulosa** cells (FSH, aromatase) → **oestrogens**.
- **Mid-cycle LH surge** → resumption of meiosis I, luteinization, ovulation; **progesterone rise** post-ovulation (corpus luteum).
- **Cervical mucus: spinnbarkeit** & ferning at ovulatory oestrogen peak—bedside correlate of Ṛtukāla.

2.3 Coordination with Āyurveda

- **Artava** is classically taught as **upadhātu of Rasa**; hence **rasa-kṣaya/āma** states distort cycles and oocyte quality (oligo-/anovulation, luteal defects).
- **Ārtavavaha srotas** integrity (uterus-vascular bed) supports **endometrial receptivity**; **Apāna-vāyu** governs cyclical egress and tubal motility.
- **Doṣa lenses:**
 - **Vāta** ↑ → anovulation, dysmenorrhoea, luteal instability.
 - **Pitta** ↑ → heavy/burning flow, luteal phase temp rise exaggerations.
 - **Kapha** ↑ → follicular arrest/PCOS phenotype, thick cervical secretions.

2.4 Applied obstetrics-gynaecology points


- **Ageing oocyte:** Aneuploidy risk rises from mid-30s; counsel **earlier conception planning** where feasible.
- **PCOS:** Hyperandrogenic, **FSH:LH imbalance** → anovulation; management aligns with **agni-dīpana, lekhana** (weight-metabolic correction) + ovulation induction when indicated.
- **Luteal phase defects:** think **rasa-rakta support** & stress modulation; correlate with **progesterone** support if clinically warranted.

3) Spermatogenesis (Śukra-nirmāṇa) — the male bija

Spermatogenesis is the continuous process of formation of male gametes (spermatozoa) from primordial germ cells (spermatogonia) in the seminiferous tubules of testes.

2. Stages of Spermatogenesis

Spermatogenesis - Stepwise Process



1. Multiplication Phase

- **Spermatogonia (diploid stem cells)** at the basal layer divide by mitosis.
- Some remain as stem cells; others differentiate into **primary spermatocytes**.

2. Growth Phase

- Primary spermatocytes (2n) enlarge and prepare for meiosis.

3. Maturation Phase

- **Meiosis I:** Primary spermatocyte → two haploid **secondary spermatocytes**.
- **Meiosis II:** Each secondary spermatocyte → two **spermatids** (haploid).
- Thus, one primary spermatocyte gives rise to **four spermatids**.

4. Spermiogenesis

- Spermatids differentiate into **spermatozoa**:
 - Nucleus condenses.
 - Acrosome forms from Golgi.
 - Flagellum develops from centriole.
 - Cytoplasm sheds.

5. Spermiation

- Mature spermatozoa released into the lumen of seminiferous tubules.



AYURVED BHARATI®

WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

www.ayurvedbharati.org

AYURVEDBHARATI.ORG



AYURVED BHARATI®

WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

www.ayurvedbharati.org

AYURVEDBHARATI.ORG



AYURVED BHARATI®

WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

www.ayurvedbharati.org

AYURVEDBHARATI.ORG



AYURVED BHARATI®

WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

www.ayurvedbharati.org

AYURVEDBHARATI.ORG




AYURVED BHARATI®

WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

www.ayurvedbharati.org

AYURVEDBHARATI.ORG





Female and Male Gametogenesis

Gametogenesis is the process of development from primordial germ cells to mature gametes: oogenesis in females and spermatogenesis in males. In both females and males, it begins with diploid germ cells that then undergo mitosis, meiosis, and cytodifferentiation into haploid gametes.

3. Duration

- Entire process: **64-72 days** in humans.
- Continuous from puberty to old age.

4. Hormonal Regulation

- **GnRH** (hypothalamus) → stimulates pituitary.
- **FSH** → acts on Sertoli cells, supports spermatogenesis.
- **LH** → stimulates Leydig cells → testosterone production.
- **Testosterone** → essential for sperm maturation and male secondary sexual characters.

5. Final Outcome

- From one primary spermatocyte: **four spermatozoa**.
- Process is **continuous, rapid, and equal**.

6. Ayurvedic Insight

- Spermatogenesis corresponds to **Shukra Dhatu formation**.
- Classical references describe continuous production of **Beeja (Shukra)** in healthy males, regulated by diet, lifestyle, and mental status.

3.1 Timeline & compartments (high-yield)

- **Site: Seminiferous tubules** (testis) — **spermatogenic epithelium** supported by **Sertoli cells**; **Leydig cells** in interstitium produce **testosterone** (LH-driven).
- **Duration: ~64-74 days** for spermatogenesis + **~12-14 days** epididymal transit/maturation.
- **Daily output: ~100-200 million** spermatozoa (wide normal).

Stages (from basement membrane to lumen):

1. **Spermatogonia (A/B)** — mitosis → pool maintenance & differentiation.
2. **Primary spermatocytes** — enter **meiosis I** → halve chromosomes.
3. **Secondary spermatocytes** — **meiosis II** → **spermatids (haploid)**.
4. **Spermiogenesis** — spermatids remodel: **acrosome**, condensed nucleus, **flagellum**, mitochondrial sheath.
5. **Spermiation** — release of spermatozoa into lumen → **epididymis** (motility & fertilizing capacity acquisition).

Blood-testis barrier (Sertoli tight junctions) protects meiotic cells from immune attack; **Androgen-binding protein (ABP)** concentrates intratubular testosterone; **Inhibin-B** from Sertoli cells exerts **FSH** feedback.

3.2 Endocrine axis & functional correlates

- **GnRH (hypothalamus)** → **LH** (Leydig → testosterone) and **FSH** (Sertoli → ABP, inhibin-B).
- **Efferent ducts** → **Epididymis** → **Vas deferens**: epididymal head-body-tail maturation (membrane remodeling,

motility).

- **Accessory glands: Seminal vesicles** (fructose, prostaglandins), **prostate** (citrate, PSA), **bulbourethral** (mucus) → final **semen** volume and buffering.

3.3 Coordination with Āyurveda

- **Śukra-dhātu** is the **culminating dhātu** (after poshana through rasa→...→majja); depletion (**śukra-kṣaya**) leads to **kārsya, klība-lakṣaṇa**, low vigour.
- **Śukravaha srotas** integrity (vr̥ṣaṇa-śukra-vāhinī pathways) maps to **testicular-epididymal** axis; **vāta** supports transport & emission, **pitta** transformation, **kapha** nutritive semen volume.
- **Doṣa pointers:**
 - **Vāta ↑**: oligo-asthenozoospermia, premature emission.
 - **Pitta ↑**: oxidative stress on sperm DNA (high ROS states).
 - **Kapha ↑**: viscous semen, poor liquefaction.

3.4 Applied andrology points

- **Heat** impairs spermatogenesis (tight clothing, saunas, fevers); the **pampiniform plexus** cools arterial blood; **varicocele** raises testicular temperature & ROS → treat when clinically significant.
- **Endocrine hits** (anabolic steroids, hyperprolactinaemia, thyroid disease) → axis suppression or poor semen quality.
- **Time-to-recovery** after gonadotoxic insults ≈ **one full spermatogenic cycle (3 months)**.

4) Head-to-Head: Oogenesis vs Spermatogenesis (exam table)

Feature	Oogenesis (Artava-nirmāṇa)	Spermatogenesis (Śukra-nirmāṇa)
When formed	Mostly fetal life ; finite pool	Post-puberty , continuous
Cell cycle arrests	Prophase I (fetal→puberty); Metaphase II (ovulation→fertilization)	None (continuous stages)
Output per cycle	Usually one secondary oocyte	Millions of sperm daily
Regulators	FSH/LH , oestrogen→LH surge → ovulation; progesterone luteal	GnRH-LH-FSH , Testosterone, Inhibin-B
Supporting cells	Granulosa/Theca , corpus luteum	Sertoli/Leydig , epididymal epithelium
Ayurvedic linkage	Ārtavavaha srotas, Apāna-vāyu, Rasa upadhātu	Śukravaha srotas, Vāta (vyāna/apāna), śukra-dhātu bala

Comparison: Oogenesis vs Spermatogenesis

Feature	Oogenesis	Spermatogenesis
Location	Ovaries	Testes (seminiferous tubules)
Onset	Fetal life	Puberty
Continuity	Discontinuous, ends at menopause	Continuous till old age
Gametes from 1 cell	1 ovum + polar bodies	4 sperms
Duration	Arrested for years, completes at fertilization	~74 days cycle
Number produced	Limited (~400 in lifetime)	Millions daily

5) Bija Quality: Ayurvedic markers & modern parallels

Āyurvedic description (concept)	Modern correlate
Śuddha Śukra: snigdha, śukla, phenodourless, compact	Normal semen: volume 1.5–6 mL, pH 7.2–8.0, rapid progressive motility, morphologically normal forms

**Āyurvedic description (concept)**

Śuddha Artava: timely, pain-light, non-fetid, non-clotty

Doṣa-duṣṭi signs (vāta/pitta/kapha)

Modern correlate

Eumenorrhoea, ovulatory cycles, normal endometrial pattern

OAT patterns, leucocytospermia/ROS, cervical mucus hostility, luteal defects

(State in answers that classical descriptors are **qualitative**, while modern lab parameters quantify similar properties.)

6) Clinical Bridges (how to use in SAQ/LAQ)

1. Preconception counselling:

- **Rasa-poshana** diet, correct **agnimāndya**, treat **āma**; sleep-stress hygiene.
- Modern: folate, iron, thyroid screen, STI screen, limit alcohol/tobacco; male antioxidant strategy if needed.

2. Infertility outline:

- **Female:** anovulation (PCOS/thyroid), tubal factors, endometrial receptivity.
- **Male:** semen analysis (2 samples), endocrine profile, rule out varicocele/infections.
- Tie back to **ārtava/śukra srotas** restoration and **doṣa-wise chikitsā** alongside indicated modern therapy.

3. ART touch-points:

- **Oocyte retrieval** follows controlled ovarian hyperstimulation (FSH/LH analogues).
- **ICSI/IVF** leverages viable **bija** even in severe OAT; still, counsel **classical svasthavṛtta** for better milieu.

7) High-Yield Numbers & Nuggets (last-minute)

- **Oocyte stock:** 6-7 million peak (fetus) → 1-2 million (birth) → 300-500k (puberty) → **~400-500 ovulations lifetime**.
- **Spermatogenesis:** **~74 days** + **~12 days** epididymis; **100-200 million/day**.
- **Ovulatory marker:** **LH surge** ~36 h before ovulation; **spinnbarkeit mucus**.
- **Male axis:** **FSH** → **Sertoli (ABP, inhibin-B)**; **LH** → **Leydig (testosterone)**.
- **Heat is harmful** to spermatogenesis; **age is harmful** to oocyte euploidy.

8) Answer Blueprints

A) LAQ (10 marks) – Describe Bija-nirmāṇa with emphasis on oogenesis and spermatogenesis; correlate with srotas and doṣa.

- Start with **Ṛtu-Kṣetra-Ambu-Bija** and **Caraka's srotas śloka** (CV 5/3).
- Oogenesis: timeline table + two-cell/two-gonadotropin model + arrests; link to **Ārtavavaha srotas** and **Apāna-vāyu**.
- Spermatogenesis: stage diagram (text) + endocrine axis + epididymal maturation; link to **Śukravaha srotas**.
- Conclude with applied infertility notes and a brief **Ayurvedic chikitsā-sūtra** (rasa-poshana, doṣa-hara).

B) SAQ (5 marks) – Enumerate key differences between oogenesis and spermatogenesis.

Use the **head-to-head table** (Section 4).

C) MCQs (single best answer)

1. Completion of **meiosis II** in the oocyte occurs:

- A) At menarche B) **After sperm entry** C) At LH surge D) At implantation

Ans: B



2. The **blood-testis barrier** is formed primarily by:
A) Leydig cells B) Peritubular myoid cells C) **Sertoli tight junctions** D) Endothelial cells
Ans: C
3. The gonadotropin that primarily **drives androgen production** in the testis is:
A) FSH B) **LH** C) Prolactin D) GH
Ans: B
4. In a normal cycle, the **mid-cycle surge** that triggers ovulation is chiefly:
A) FSH B) **LH** C) Progesterone D) Prolactin
Ans: B
5. **Artava** is classically aligned with which dhātu relationship?
A) Māṃsa upadhātu B) **Rasa upadhātu** C) Meda upadhātu D) Majjā upadhātu
Ans: B

9) References

Classical (primary authorities)

- **Caraka Saṃhitā, Vimāna-sthāna 5 (Srotovimāna)** — definition and doctrinal role of **srotas** (quoted above: 5/3); mūla-descriptions for reproductive srotas used for correlation in answers.
- **Suśruta Saṃhitā, Śārīra-sthāna** — garbha-sambhava framework (Ṛtu-Kṣetra-Ambu-Bīja) and reproductive organ descriptions supporting **Ārtavavaha/Sukravaha** pathways.
- **Aṣṭāṅga Hṛdaya, Śārīra-sthāna** — consolidated teachings on reproductive physiology; phases around **Ṛtukāla** and qualities of **śuddha śukra/artava** in commentarial traditions.

Note: Only a **verified śloka** (CV 5/3) is quoted in Devanāgarī to maintain textual accuracy as per your instruction. All other classical points are aligned with the above sthānas without risking unverified meter.

Modern & standard texts

- **Williams Obstetrics** (ovarian cycle, oogenesis, endocrine control).
- **Guyton & Hall / Ganong** (hypothalamo-pituitary-gonadal axis).
- **Campbell-Walsh Urology / Andrology reviews** (spermatogenesis, epididymal maturation, semen parameters).
- **WHO Laboratory Manual for the Examination and Processing of Human Semen** (latest edition) — benchmarks for semen quality.
- **Dutta's Gynecology** — menstrual physiology, ovulation indicators, clinical correlations.

10) Quick Recap (30-second lock-in)

- **Bīja-nirmāṇa = gamete formation** carried by **srotas**; cite **CV 5/3**.
- **Oogenesis:** fetal origin, **two arrests**, one oocyte per cycle; governed by **FSH/LH**, granulosa-theca teamwork.
- **Spermatogenesis:** continuous post-puberty, **~3 months** cycle, Sertoli/Leydig axis, epididymal finishing.
- **Āyurveda: Ārtava = Rasa-upadhātu; Śukra-dhātu** culmination; integrity of **Ārtava/Sukra-vaha srotas + doṣa-sāmyā** underpins fertility.