Unit 4 — Sañjñāharaṇa in Netraroga (Anesthesia in Ophthalmology)

Unit 4 — Sañjñāharaṇa in Netraroga (Anesthesia in Ophthalmology)

Goal: by the end of this chapter you will be able to choose, administer, and monitor appropriate anesthesia for common ophthalmic procedures, anticipate complications (especially the oculocardiac reflex), and write exam-ready notes that integrate Ayurvedic context with contemporary standards of care.

1) Concept & Scope

Sañjñāharaṇa (abolition of sensation/awareness for painless procedures) is indispensable in Śālākya—Netra practice. While classical Ayurveda acknowledges the need to reduce pain, fear, and reflexes before sharp/operative measures, today's ophthalmic anesthesia is delivered by local/topical techniques, regional orbital blocks, monitored sedation, and general anesthesia—selected according to surgery type, patient factors, and surgeon skill. Your exam answers should show:

- Clear **indications/contraindications** for each technique.
- Drug names, concentrations, volumes, and adjuvants.
- Complication recognition + first-line management (especially LAST, retrobulbar hemorrhage, globe perforation, brainstem anesthesia, oculocardiac reflex).

2) Pre-anesthetic Assessment & Preparation (Netra Focus)

History & risks: previous reactions to local anesthetics, bleeding disorders/anticoagulants, uncontrolled **HTN/DM**, asthma/COPD, **axial high myopia** (globe perforation risk), glaucoma, psychiatric illness, communication barriers.

Examination: vitals, airway (if sedation/GA), ocular status (axial length if available), **IOP trends**, extra-ocular movements, lids, conjunctiva.

Consent & instructions: fasting (if sedation/GA), avoid eye rubbing, arrange escort.

Monitoring (minimum): pulse oximetry, NIBP, +/- ECG in elderly/comorbid.

Rescue readiness: oxygen, suction, bag-mask, IV access, **lipid emulsion 20%**, benzodiazepine for seizures, **atropine** for bradycardia, emergency cart.

3) Types of Ophthalmic Anesthesia

A) Topical Anesthesia (surface corneal/conjunctival block)

Agents (typical):

- **Proparacaine 0.5%** (rapid onset ~20–30 s; duration ~10–15 min)
- **Tetracaine 0.5%** (similar profile; more sting)
- Lignocaine (lidocaine) 2% gel (longer contact, useful in phaco)
- Oxybuprocaine 0.4% (where available)

Indications: phacoemulsification (many cases), pterygium excision (with subconjunctival supplement), foreign-body removal, suture removal, diagnostic procedures.

Advantages: easy, guick, maintains ocular motility/airway reflexes.

Limitations: no akinesia, poor for long/painful intraocular steps; epithelial toxicity if repeatedly instilled.

[©] Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.



Pearls: add intracameral preservative-free lidocaine 1% (0.1-0.2 ml) during cataract as needed.

B) Infiltration & Regional Anesthesia

1) Subconjunctival Anesthesia

- Drug: lignocaine 2% (0.2-0.5 ml) +/- small adrenaline for lid hemostasis (avoid intraocular vasoconstrictor).
- **Use:** pterygium, conjunctival suturing, superficial procedures.

2) Intracameral Anesthesia

- **Drug: preservative-free** lignocaine 1% (0.1-0.2 ml) via paracentesis.
- **Use:** cataract (with topical), anterior chamber manipulation.

3) Sub-Tenon's (Episcleral) Block — preferred regional for many intraocular surgeries

- Cannula: blunt 19-23G through a small inferonasal conjunctival/Tenon's nick.
- Volume/Drugs: 3-5 ml mixture (e.g., lignocaine 2% + bupivacaine 0.5%), often with hyaluronidase 7.5-15 IU/ml.
- Advantages: excellent anesthesia, good akinesia with low risk of globe perforation, optic nerve injury, or brainstem spread.
- Indications: cataract, trabeculectomy, many vitreo-retinal procedures in cooperative adults.

4) Peribulbar Block

- **Needle:** 25–27G, 25–38 mm; 1–2 injection sites (inferotemporal ± superonasal).
- Volume/Drugs: 6-10 ml of lignocaine 2% + bupivacaine 0.5% + hyaluronidase 7.5-15 IU/ml; onset 5-10 min.
- Advantages: reliable akinesia; safer than retrobulbar.
- Cautions: chemosis/eyelid edema, raised IOP; gentle ocular compression (e.g., Honan device 10-15 mmHg for ~10 min) can help.

5) Retrobulbar Block (posterior intraconal) — declining use

- Needle: 23-25G, ~31-38 mm via inferotemporal fornix; **2-4 ml** of agent.
- Pros: rapid, dense block.
- Cons/risks: retrobulbar hemorrhage, optic nerve injury, globe perforation, brainstem anesthesia; now replaced by sub-Tenon's/peribulbar in many centers.

6) Facial Nerve Blocks (for lid akinesia in selected lid/orbital surgery)

- Van Lint, O'Brien, Atkinson techniques—targeting branches of VII nerve to prevent forceful squeezing.
- Often unnecessary if peribulbar/sub-Tenon's already provide adequate orbicularis relaxation; still useful in blepharoplasty, entropion/ectropion repair.

7) Eyelid Infiltration

- Drug: lignocaine 2% with or without adrenaline 1:200,000 for hemostasis (avoid if peripheral perfusion is compromised).
- **Use: chalazion**, lid lacerations, minor lid procedures; inject **subcutaneously along lid margin**, respecting septum and avoiding globe.

C) Monitored Anesthesia Care (MAC) / Sedation

- Goals: anxiolysis and analgesia without airway compromise.
- Common regimens:

[©] Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.



- Midazolam 0.5-2 mg IV titrated;
- Fentanyl 25-50 μg IV (cautious in elderly);
- Dexmedetomidine infusion for cooperative sedation (bradycardia risk—monitor).
- Precautions: supplemental oxygen via nasal cannula; continuous SpO₂/NIBP; avoid oversedation (especially in topical cases where patient feedback is valuable).

D) General Anesthesia (GA)

Indications: infants/children, uncooperative/anxious adults, long vitreoretinal/orbital surgeries, open globe trauma, combined procedures.

Airway/Agents: IV induction (e.g., **propofol**), **sevoflurane** maintenance; **LMA** or **ETT** based on aspiration risk; multimodal analgesia; antiemetic prophylaxis (**ondansetron**).

Special issues: prevention/management of **oculocardiac reflex**, gentle emergence to avoid IOP spikes, protection of the operated eye.

4) Drugs in Ophthalmic Anesthesia — Quick Reference

Class	Examples	Key points (dose/conc.)	
Local anesthetics	Lignocaine 2%, Bupivacaine 0.5%, Ropivacaine 0.5-0.75%	Onset: lignocaine fast; duration: bupivacaine long. Max safe dose (general guide): lignocaine 3-5 mg/kg (plain), up to ~7 mg/kg with adrenaline; bupivacaine ≤ 2.5 mg/kg; ropivacaine ≤ 3 mg/kg. Use small orbital volumes.	
Topicals	Proparacaine 0.5%, Tetracaine 0.5%, Lignocaine gel 2%	Rapid onset; avoid frequent repeat → epithelial toxicity/delayed healing.	
Adjuvants	Hyaluronidase 7.5-15 IU/ml	Improves spread/akinesia; beware allergy.	
Vasoconstrictor (limited use)	Adrenaline 1:200,000	For skin/lid infiltration to reduce bleeding; avoid mixing in peri/retrobulbar solution (ischemia risk).	
Sedatives/analgesics	Midazolam, Fentanyl, Dexmedetomidine	Titrate slowly; monitor airway.	
Anticholinergic	Atropine (OCR)	For bradycardia during extraocular muscle traction.	
Rescue for LAST	Lipid emulsion 20%	Bolus 1.5 ml/kg , then infusion per protocol if systemic toxicity occurs.	

5) Complications & Their Management

A) Oculocardiac Reflex (OCR)

Trigger: traction on extraocular muscles, pressure on globe, retrobulbar injection.

Pathway: Afferent—V1 (trigeminal) → brainstem → Efferent—vagus → bradycardia, AV block, asystole in severe cases

Management: Stop stimulus, call out; ensure oxygenation, deepen anesthesia/sedation; **atropine IV** if bradycardia persists. **Prevention:** adequate block; prophylactic anticholinergics in high-risk pediatric strabismus surgery.

B) Retrobulbar Hemorrhage

Clues: sudden proptosis, tense lids, pain, rapidly rising IOP, loss of red reflex.

Immediate actions: stop, call help, IOP-lowering measures (IV mannitol if available), lateral canthotomy/cantholysis if optic nerve perfusion is threatened; postpone surgery.

[©] Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.

C) Globe Perforation

Risk: high axial myopia, posterior staphyloma, long sharp needle use. **Signs:** sudden pain, hypotony, decreased vision, vitreous hemorrhage.

Action: stop injection, urgent retina consult, B-scan/ocular exam, manage retinal tears/detachment promptly.

D) Local Anesthetic Systemic Toxicity (LAST)

Early signs: perioral numbness, tinnitus, metallic taste, agitation → seizures, CNS depression; later arrhythmias.

Treatment: secure airway, oxygen, benzodiazepine for seizures, start 20% lipid emulsion (bolus 1.5 ml/kg → infusion), ALS protocol.

E) Brainstem Anesthesia

Mechanism: inadvertent spread along optic nerve sheath. **Presentation:** contralateral amaurosis, dysarthria, apnea.

Management: airway support, oxygenation, cardiovascular stabilization; observe till recovery.

F) Myotoxicity / Diplopia

Cause: direct injection into extraocular muscle, high-concentration bupivacaine. **Prevention:** correct needle path; prefer **sub-Tenon's** or lower-concentration mixes.

G) Topical Toxicity

Excess proparacaine/tetracaine → punctate keratopathy, delayed healing; counsel against self-use.

6) Choosing the Technique — Procedure-wise Matrix

Procedure	First choice	Alternatives	Notes
Phacoemulsification (adult)	Topical + intracameral	Sub-Tenon's / Peribulbar (anxious/unsteady)	No globe akinesia with topical; talk through steps.
Trabeculectomy	Sub-Tenon's	Peribulbar	Avoid adrenaline near filtration tissues.
Pterygium	Topical + subconjunctival	Sub-Tenon's for extensive cases	Adequate topical is often sufficient.
Vitreoretinal (selected)	Sub-Tenon's/Peribulbar	GA (long/complex)	Surgeon preference; patient cooperation crucial.
Strabismus (children)	GA	_	OCR risk high; use anticholinergic prophylaxis.
Chalazion / Lid surgery	Lid infiltration	+ Sedation if anxious	Consider facial nerve block for blepharoplasty.
Open-globe trauma	GA	_	Prevent extrusion; do not compress globe.

7) Practical Steps (checklist style)

- 1. Confirm plan with surgeon (topical vs block vs GA).
- 2. Mark eye, establish IV line (for blocks/MAC/GA), monitors on.
- 3. **Asepsis:** povidone-iodine 5% into fornices before block.
- 4. **Perform block** (prefer **sub-Tenon's** where feasible).
- 5. **Ocular compression** (peribulbar) to reduce IOP and spread agent.
- 6. Reassess akinesia/analgesia before draping; top-up if needed.

[©] Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.



- 7. Intra-op vigilance: OCR watch, patient communication (topical).
- 8. **Post-op:** shield, analgesia, nausea control, discharge advice.

8) Ayurvedic Context (framing for viva)

In classical surgical thought, eliminating pain and fear before **śastra/yantra** use is emphasized; in the **ūrdhvajatru** region this aligns with **gentle handling, mind-body calming**, and appropriate **śodhana/śamana** sequencing for recovery. In practice today, **contemporary ophthalmic anesthesia** is the standard of care, while **pathya-apathya** (light diet, sleep hygiene, wind/sun protection) and **snehana/rasāyana after** the acute phase may aid ocular surface homeostasis. Use this framing to show integrative understanding—**do not substitute modern anesthesia with traditional measures in the operative setting.**

Assessment (Exam-Ready)

Long Essays (10 marks each; attempt any 1)

- 1. Enumerate the types of anesthesia used in ophthalmology and detail indications, drugs (concentration/volume), steps, and complications of sub-Tenon's and peribulbar blocks.
- 2. **Discuss the oculocardiac reflex** in detail—mechanism, risk factors, prevention, and stepwise management. Add how choice of block modifies its incidence.

Short Essays (5 marks each; attempt any 3)

- 1. Compare topical anesthesia vs intracameral anesthesia for cataract surgery.
- 2. Outline the technique, advantages, and risks of a **retrobulbar block** and explain why many centers prefer **sub-Tenon's** today.
- 3. Write notes on **sedation (MAC)** in ophthalmology—drug choices and safety.
- 4. Indications and technique of facial nerve blocks in eyelid surgery.

Short Notes (3 marks each; attempt any 4)

- Hyaluronidase: role, dose, cautions.
- Lipid emulsion therapy in LAST (dose and triggers).
- Retrobulbar hemorrhage: recognition and immediate actions.
- Intracameral lignocaine: dose, use, precautions.
- Adrenaline in lid infiltration: when to avoid.
- Brainstem anesthesia: presentation and first aid.

MCQs (1 mark \times 5)

- 1. Typical hyaluronidase concentration in orbital blocks is:
 - a) 1-2 IU/ml b) **7.5-15 IU/ml** c) 50 IU/ml d) 100 IU/ml
- 2. First-line response to **oculocardiac reflex** is:
 - a) Give atropine immediately b) Increase IV fluids c) **Stop the stimulus and ensure oxygenation** d) Start chest compressions
- 3. The **safest regional technique** for many intraocular surgeries with lowest risk of globe perforation is:
 - a) Retrobulbar b) **Sub-Tenon's** c) Peribulbar d) Facial nerve block
- 4. Intracameral lignocaine for phaco is usually given as:
 - a) 1 ml of 2% b) **0.1-0.2 ml of 1% (preservative-free)** c) 1 ml of 1% with adrenaline d) 2 ml of 0.5%
- 5. The afferent limb of the oculocardiac reflex is via:
 - a) Facial nerve b) Optic nerve c) Trigeminal nerve (V) d) Glossopharyngeal nerve

Answer key: 1-b, 2-c, 3-b, 4-b, 5-c.

[©] Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only. Unauthorized reproduction, distribution, or commercial use is strictly prohibited.





60-Second Recap (for viva)

- **Topical + intracameral** is enough for many phaco cases; **sub-Tenon's** is the versatile, safer regional block; **peribulbar** gives strong akinesia; **retrobulbar** is powerful but riskier.
- Know **drug concentrations, volumes, adjuvants (hyaluronidase)**, and **what not to mix** (avoid adrenaline in orbital solutions).
- Recognize and manage **OCR**, **retrobulbar hemorrhage**, **LAST**, **globe perforation**, **brainstem anesthesia**—these fetch marks.
- Integrate Ayurvedic framing ethically: pain-free, calm, and protected eyes—but use modern anesthesia standards in the OR.

[©] Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.