



Chapter 6. Handling Medical Emergencies Part 1. Respiratory Emergencies

Chapter 6: Handling Medical Emergencies

Part 1: Respiratory Emergencies

(Managing choking, asthma attacks, and allergic reactions effectively)

1 What Are Respiratory Emergencies?

Respiratory emergencies involve **difficulty or inability to breathe** due to blocked airways, lung conditions, or severe allergic reactions. These conditions can become life-threatening within minutes if not managed properly.

2 Choking (Airway Obstruction)

What Happens?

Choking occurs when the airway is **partially or completely blocked**, usually by food, vomit, or an object.

How to Recognize It:

Partial Obstruction	Complete Obstruction
Coughing forcefully	Unable to speak, breathe, or cough
Gasping	Silent attempts to breathe
Noisy breathing	Clutching the throat (universal sign)

First Aid for Choking (Adults & Children >1 Year)

A. If the person is coughing strongly:

- Encourage coughing.
- Do not hit their back or interfere.

B. If airway is blocked and they cannot breathe or speak:

1. **Ask: "Are you choking?"**
2. **Call 112 immediately.**
3. **Give 5 back blows:**
 - Stand behind, support chest, lean person forward.
 - Hit firmly between the shoulder blades with heel of your hand.
4. **Give 5 abdominal thrusts (Heimlich Maneuver):**
 - Stand behind, place fist above navel, other hand over it.
 - Pull inwards and upwards.
5. **Alternate 5 back blows + 5 thrusts** until object comes out or they collapse.

If unconscious, start **CPR** and check mouth for visible object during breaths.



For Infants (<1 Year)

- **Never use abdominal thrusts.**
- Place baby face down on your forearm, head lower than chest.
- Give **5 gentle back blows** between shoulder blades.
- Turn face-up and give **5 chest thrusts** using 2 fingers (like CPR but slower).
- Repeat until breathing returns or CPR is needed.

3 Asthma Attack

What is Asthma?

A **chronic lung condition** where airways become inflamed, narrow, and produce excess mucus. Triggers include dust, cold, exercise, smoke, and infections.

Signs of an Asthma Attack:

- Wheezing or whistling sound when breathing
- Shortness of breath
- Chest tightness
- Coughing (especially at night)
- Use of accessory muscles to breathe
- Speaking in broken sentences

First Aid for Asthma

1. **Keep the person calm and sitting upright.**
2. **Assist with their inhaler (usually a blue “reliever” like salbutamol):**
 - Shake inhaler
 - Attach spacer (if available)
 - Give **1 puff every 30-60 seconds**, up to 10 puffs
3. **Call 112 if:**
 - No relief after 10 puffs
 - Breathing worsens
 - Person becomes drowsy, pale, or unresponsive

Using a Metered-Dose Inhaler (MDI) Correctly

With Spacer	Without Spacer
Shake inhaler	Shake inhaler
Insert into spacer	Hold upright
Inhale slowly through mouthpiece	Exhale fully, then inhale deeply while pressing canister
Hold breath 5-10 sec	Hold breath 5-10 sec
Wait 30-60 sec before next puff	Same

4 Anaphylaxis (Severe Allergic Reaction)

What Is It?

An **extreme allergic reaction** that causes rapid airway swelling, low blood pressure, and can be fatal if untreated.

Common Triggers:

- Food (nuts, shellfish, eggs)
- Insect stings
- Medications (e.g., penicillin)
- Latex

Symptoms of Anaphylaxis:

- Swelling of lips, tongue, or throat
- Difficulty breathing or wheezing
- Rash or hives
- Nausea, vomiting
- Feeling dizzy or faint
- Rapid pulse, low BP
- Sense of doom or confusion

First Aid for Anaphylaxis

1. **Call 112 immediately.**
2. **Lay the person flat** (unless breathing worsens—then sit).
3. **Give Epinephrine auto-injector (e.g., EpiPen):**
 - Inject into outer mid-thigh (even through clothes).
 - Hold for 3–5 seconds.
4. **Repeat dose after 5 minutes if no improvement and second pen is available.**
5. Monitor breathing and be ready to give CPR if needed.

Do not give food, water, or medication by mouth.

How to Use an Epinephrine Auto-Injector

1. Remove safety cap.
2. Firmly press into mid-thigh at 90°.
3. Hold for 3–5 seconds.
4. Remove and massage area.
5. Call EMS even if symptoms improve.

5 Self-Check Questions

1. What is the first step when you see someone choking?
2. How many puffs of an asthma reliever can be given in one attack?
3. Name two signs of anaphylaxis.



4. Where is an EpiPen injected?
 5. Why is it important to still call EMS after using an auto-injector?
1. Ask if they are choking and encourage coughing if possible.
 2. Up to 10 puffs, 30–60 seconds apart.
 3. Swelling of tongue/throat, wheezing, rash, dizziness.
 4. Into the outer thigh.
 5. Because symptoms can return; further medical treatment is needed.

Key Take-Home Points

- Choking requires **back blows and abdominal thrusts**—know how to act fast.
 - Asthma is common and often manageable with **quick-relief inhalers**.
 - Anaphylaxis is a medical emergency—**epinephrine and EMS are lifesaving**.
 - Practising how to use inhalers and EpiPens can prepare you for real emergencies.
-