

#### WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

## vi. Purification of bioactive compounds through various chromatographic methods

## **Context and Rationale for Purification**

#### 1. Significance of Bioactive Isolation

- Traditional polyherbal formulations in Ayurveda feature multiple compounds. Isolating specific bioactive molecules (alkaloids, flavonoids, terpenoids, glycosides) can help:
  - 1. Confirm pharmacological activities in mechanistic or clinical studies,
  - 2. Standardised herbal extracts for consistent potency,
  - 3. **Develop** advanced formulations or novel drug leads.
- o Purified reference standards also facilitate quality control and authenticity verification.

### 2. Complexity of Botanical Matrices

- Raw plant extracts often contain dozens to hundreds of phytochemicals with varying polarity, molecular weight, and thermal stability.
- Effective purification strategies rely on stepwise fractionation to separate compounds with distinct physicochemical traits.

## 3. Integration With Other Methods

Chromatographic purification is frequently **preceded** by broad extraction techniques (maceration, Soxhlet, supercritical CO<sub>2</sub> extraction) and **followed** by confirmatory analysis (spectroscopy, mass spectrometry, NMR) to verify purity and structure.

# **Chromatographic Principles**

#### 1. Modes of Chromatography

- **Adsorption Chromatography**: Solute molecules adsorb onto a solid stationary phase; differential interactions partition compounds.
- **Partition Chromatography**: Molecules distribute between a stationary phase (often a bonded liquid on a support) and the mobile phase.
- **Ion-Exchange Chromatography**: Separation based on charge interactions between analytes and a charged stationary phase.
- Size-Exclusion (Gel Filtration): Separates molecules by size/shape, with larger molecules eluting earlier than smaller ones.

#### 2. Mobile and Stationary Phases

- Typically a solid or immobilized phase (silica, polymer resin, reversed-phase C18, etc.) and a mobile phase (liquid or gas).
- Polarity, pH, ionic strength, and temperature influence compound retention or elution.

#### 3. Elution Strategies

- Isocratic (constant solvent composition) or gradient (varying solvent polarity over time).
- In plant extract purification, gradient elution is common to systematically separate compounds of increasing polarity or partition coefficients.

## **Chromatographic Methods for Bioactive Purification**

## Column Chromatography (Open-Column or Gravity-Driven)

## 1. Fundamental Setup

- A glass or plastic column packed with a stationary phase (e.g., silica gel, alumina) over which plant extract in a suitable solvent is loaded.
- o Elution proceeds by adding a mobile phase (often increasing polarity in stepwise manner).

## 2. Applications

- Widely used as a **preparative** method for fractionation in the lab, or as a simpler approach to segregate broad classes (non-polar vs. polar fractions).
- Cost-effective but relatively slow, requires large volumes of solvent, and fraction collection/monitoring by TLC or UV detection.

<sup>©</sup> Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.



#### WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

### 3. Flash Chromatography

- A more modern variant of column chromatography using pressurized gas or pumps for faster flow, smaller particle size stationary phases.
- o Improves resolution and speed, widely used for medium-scale isolation of desired compounds.

## **High-Performance Liquid Chromatography (HPLC)**

#### 1. Principle and Hardware

- **High-pressure pumps** drive the mobile phase through a column packed with small, high-efficiency particles (C18 reversed-phase, normal-phase silica, etc.).
- o UV-Vis, diode-array (DAD), or MS detectors track eluting peaks in real-time, facilitating fraction collection.

#### 2. Reversed-Phase HPLC

- Most common for semi-polar or polar compounds from herbal extracts.
- Gradual gradient (e.g., water with 0.1% formic acid to acetonitrile) yields good separation of complex mixtures
- o Fractions corresponding to peaks of interest are collected, typically repeated to improve purity.

## 3. Preparative HPLC

- Larger diameter columns (semi-preparative or prep scale) with higher flow rates for producing multi-gram quantities of purified phytoconstituents.
- Balances throughput, resolution, and cost. Often used in advanced R&D labs or herbal manufacturing settings needing high-purity active principles.

## Thin-Layer Chromatography (TLC)/High-Performance Thin-Layer Chromatography (HPTLC)

#### 1. Basic TLC

- **Stationary phase**: Silica or reversed-phase plates. The sample is spotted on the plate; mobile phase climbs by capillary action, separating compounds by differential adsorption.
- Qualitative or semi-quantitative method for **fingerprinting**, verifying fraction composition, or quick check of extraction progress.

## 2. HPTLC

- More refined version with smaller particle size, automated sample application, and densitometric scanning.
- Enhanced resolution, reproducibility, and the possibility to identify multiple components rapidly, useful in routine QA for polyherbal mixtures.
- Not typically used for large-scale purification but excellent for **monitoring** fraction purity or standardization.

## **Gas Chromatography (GC)**

## 1. Suitability

- Applied to **volatiles** (essential oils, low molecular weight terpenes) or derivatives.
- The plant extract often needs derivatization if compounds aren't volatile or stable enough for GC conditions.
- Highly relevant for aromatic plants (cinnamon, clove, fennel) in analyzing essential oil composition, but not for typical large, polar, or heat-labile molecules.

## 2. Advantages

- High resolution, robust library-based identification (mass spectra).
- Fast analysis, widely used for flavor/fragrance compounds or small, thermally stable constituents.

## Ion-Exchange and Size-Exclusion Chromatography

#### 1. Ion-Exchange

- o Targets charged compounds (alkaloid salts, ionic polysaccharides).
- o Tuning pH and ionic strength for stepwise elution.

### 2. Size-Exclusion (SEC/Gel Filtration)

- Useful for isolating proteins, large polysaccharides (e.g., mucilages, starch modifications) from smaller phenolics.
- Rarely used for typical small-molecule phytochemicals but essential for specialized Ayurvedic formulations requiring macromolecule fractionation.

<sup>©</sup> Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only Unauthorized reproduction, distribution, or commercial use is strictly prohibited.



#### WHERE CLASSICAL WISDOM MEETS INTELLIGENT LEARNING

# **Considerations for Scale-Up and Quality Assurance**

## **Process Optimization**

#### 1. Solvent Recycling

- Large-scale extraction can be solvent-intensive. Installing solvent recovery systems (distillation, condensation) reduces cost and environmental impact.
- Ensuring minimal residual solvent in the final product is crucial for safety, especially if using non-food-grade solvents.

#### 2. Automation and Monitoring

- Automated fraction collectors, online UV or MS detection, use of advanced software controlling gradient pumps enhance reproducibility.
- Real-time data capture allows immediate decisions on fraction pooling or re-running partially resolved fractions.

## **Regulatory and Safety Issues**

#### 1. Residual Solvents

- Strict limits set by pharmacopoeial standards (e.g., ICH guidelines for solvent residues in pharmaceuticals).
- Ethanol or water-based processes are generally more consumer-friendly.

#### 2. Heavy Metal and Microbial Load

- o Even after compound isolation, verifying no contamination from raw materials or process steps.
- GMP guidelines require validated cleaning procedures for columns, equipment to prevent crosscontamination between batches.

#### 3. Authenticity and Traceability

- o Standard operating procedures ensuring each extracted batch ties back to raw material batch codes.
- Transparent documentation fosters consumer trust, helps meet AYUSH, FDA, or EU herbal directive compliance.

# **Concluding Remarks**

**Chromatographic purification** of bioactive compounds from medicinal plants is central to **modern herbal R&D**, bridging:

- 1. Traditional knowledge about multi-ingredient synergy (as in Ayurveda) with
- 2. **Scientific demands** for reproducible, high-purity fractions to confirm **pharmacological activity** and ensure **regulatory compliance**.

Techniques like **Column Chromatography**, **Soxhlet**, **Flash Chromatography**, **HPLC**, **HPTLC**, and **Supercritical Fluid Extraction**—each with distinct principles and hardware—enrich the researcher's toolkit for systematically isolating target molecules, verifying authenticity, and scaling production. Leveraging **polarity-based** fractionation strategies and **appropriate solvents** ensures that essential constituents are selectively recovered with minimal degradation.

Ultimately, a well-conceived **chromatographic workflow**—aligned with good manufacturing practices, ecological responsibility, and ethical sourcing—amplifies the potential of **Ayurvedic** and other herbal resources to yield safe, consistent, and efficacious products for global health applications.

<sup>©</sup> Ayurvite Wellness Pvt Ltd. All rights reserved. This PDF is for personal use only. Unauthorized reproduction, distribution, or commercial use is strictly prohibited.